





# Healthier Communities... Healthier Families

## Introducing a New Standard for Improved Quality of Life in Montana!

Montana's New <u>Clean Indoor Air Act</u> provides citizens with protection from exposure to secondhand tobacco smoke in the workplace, and in public places such as restaurants, shopping areas, schools, bowling alleys, and some bars. The result? Healthier communities and families in Montana.

### Secondhand Smoke Causes Cancer and Makes Many People Who Don't Smoke Sick. The New Law Will Offer Protection.

- Secondhand smoke is a mixture of the smoke given off by the burning end of a cigarette or other tobacco product and the smoke exhaled by smokers.<sup>1,2</sup>
- Secondhand smoke is hazardous and contains a mixture of more than 4,000 chemicals, at least 50 of which are known or probable human carcinogens. Breathing it causes CANCER.<sup>3,4</sup>
- The U.S. Surgeon General has concluded that exposure to secondhand smoke is a common public health hazard that is *completely preventable*. Exposure can be dramatically reduced by eliminating smoking in all enclosed public places and workplaces. <sup>5, 6, 7, 8</sup>

#### Montana's new law will help to reduce needless death.

- Each year, an estimated 175 Montanans who never themselves smoked, die from breathing someone else's tobacco smoke in the workplace or home.<sup>9</sup>
- Each year in the U.S., secondhand smoke is associated with up to 62,000 heart disease deaths and 3,000 lung cancer deaths.<sup>1</sup>

#### **Solution** Montana's new law will help to improve the health of citizens.

- Breathing secondhand smoke is associated with an increased risk for sudden infant death syndrome (SIDS), asthma, bronchitis, and pneumonia in children.<sup>1, 10</sup>
- Each year, secondhand smoke is associated with 8,000 26,000 new asthma cases in children and an estimated 150,000 300,000 new cases of bronchitis and pneumonia in children less than 18 months old (7,500 15,000 of which require hospitalization).  $^{4,11,12}$
- A study in Helena, Montana found that the number of heart attack victims admitted to the regional hospital dropped by nearly 60% during the first 6 months of a new smokefree community ordinance taking effect.<sup>13</sup>
- Regular exposure to secondhand smoke, such as in restaurants and bars, increases one's chance of stroke by as much as 50 percent.<sup>14</sup>
- Not only will Montana's new law reduce exposure to secondhand smoke among workers and the public, it will help smokers quit and/or reduce the number of cigarettes they smoke and will provide a stronger positive behavioral influence for children.<sup>15</sup>

#### Montana's new law will allow citizens to participate more actively and fully in their communities.

- The short-term effects of breathing secondhand smoke such as wheezing, asthma attacks, shortness of breath, and excessive coughing – last for hours, and sometimes days after exposure. 16
- As little as 30 minutes of exposure to secondhand smoke puts certain individuals at greater risk of heart attack. The Centers for Disease Control (CDC) warns that people with known heart disease should avoid all indoor environments that permit smoking.<sup>16</sup>

#### Montana's new law extends important protection to workers who traditionally have been exposed to high levels of secondhand smoke.

- Secondhand smoke in bars and casinos is 3 to 6 times more concentrated than in other smoke-filled workplaces. Secondhand smoke levels in restaurants are nearly TWICE those found in smoke-filled offices. 17
- Food service workers have a 50% greater risk of dying from lung cancer than the general population, in part, because of secondhand smoke exposure in the workplace. 18, 19
- Breathing the air for just two hours in a smoky bar produces similar health effects as smoking FOUR cigarettes.<sup>20</sup>
- Studies have shown marked improvement in bar workers' health within the first few months of implementing a smokefree policy.<sup>21</sup>

#### Montana communities & citizens strongly support a statewide smokefree law.

■ 71% (nearly three-fourths) of Montanans polled support the statewide smokefree public places law.<sup>22</sup>

#### Need More Information on Montana's New Clean Indoor Air Act?

Contact the Montana Tobacco Use Prevention Program for more information and for the names of local educators or representatives who may be able to answer your questions.

#### **Montana Tobacco Use Prevention Program**

PO Box 202951

Helena, MT 59620-2951

Toll-Free Information Line: 866-787-5AIR (866-787-5247)

- 1. National Cancer Institute. Health Effects of Exposure to Environment Tobacco Smoke. Smoking and Tobacco Control Monograph No. 10. Bethesda, MD: U.S. Department of Health and Human Services, National Institutes of Health, National Cancer Institute; 1999. NIH Pub. No. 99-4645.
- 2. National Toxicology Program. 10th Report on Carcinogens. Research Triangle Park. NC: U.S. Department of Health and Human Services, Public Health Service, National Toxicology Program. Dec 2002
- 3. Pirkle JL, Flegal KM, Bernert JT, Brody DJ, Etzel RA, Maurer KR. Exposure of the U.S. population to environmental tobacco smoke: The Third National Health and Nutrition Examination Survey, 1988 to 1991. Journal of the
- 4. U.S. Environmental Protection Agency. Respiratory Health Effects of Passive Smoking: Lung Cancer and Other Disorders. Washington, DC: U.S. Environmental Protection Agency; 1992
- 5. Hopkins DP, et., al. Reviews of evidence regarding interventions to reduce tobacco use and exposure to environmental tobacco smoke. Am J Prey Med 2001;20 (2 suppl): 16-66.
- 6. Heloma A, Jaakkola MS. Four-year follow-up of exposure, attitudes and smoking behaviour following enactment of Finland's national smoke-free work-place law. Addiction 2003;98: 1111-7.
- 7. An air quality survey of respirable particles and particulate carcinogens in Delaware hospitality venues before and after a smoking ban. A report for the Delaware Tobacco Prevention Coalition, Feb 2003. Silver Springs, MD. 8. Travers M, Cummings KM, Repace J, Hyland A. Indoor air quality before and after the New York state clean indoor air law in western New York hospitality venues, July-August 2003. Paper presented at Society for Research on Nicotine and Tobacco Annual Conference, Scottsdale, Arizona, February 18-20, 2004.
- 9. Centers for Disease Control and Prevention (CDC), National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2002. Tobacco Control State Highlights 2002: Impact and
- 10. U.S. Department of Health and Human Services, Public Health Service, Office of the Surgeon General. Rockville, MD: U.S. Department of Health and Human Services, Public Health Service, Office of the Surgeon
- 11. CDC. 2nd National Report on Human Exposure to Environmental Chemicals: Tobacco Smoke. U.S. Dept. of Health and Human Services, CDC, National Center for Environmental Health; 2003:80. NCEH Pub No. 03-0022. 12. CDC. State-specific prevalence of cigarette smoking among adults, and children's and adolescents' exposure to environmental tobacco smoke—United States, 1996. Morbidity and Mortality Weekly Report 1997;
- 13. Sargent, RP, Shephard, Rs, Glantz, SA. Immediate Reduction in Acute Myocardial Infarctions After the Implementation of a Comprehensive Smokefree Ordinance: American College of Cardiology, 52nd Scientific Sessions, Chicago, IL. April 2003.
- 14. Zhang, X. et al., "Association of Passive Smoking by Husbands with Prevalence of Stroke among Chinese Women Nonsmokers," American Journal of Epidemiology, 2005; 161(3): 213-218.

  15. U. S. Department of Health and Human Services, Reducing Tobacco Use: A Report of the Surgeon General. Atlanta, Georgia: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention,
- National Center for Chronic Disease Prevention and Health Promotion, 2000. 16. Otsuka R, Watanabe H, Hirata K, Tokai K, Muro T, Yoshiyama M, Takeuchi K, Yoshikawa J., Acute effects of passive smoking on the coronary circulation in healthy young adults; JAMA. 2001 Jul 25;286(4):436-41.
- 17. Shopland, D.R.; Anderson, C.M.; Burns, D.M.; Gerlach, K.K., "Disparities in smoke-free workplace policies among food service workers," Journal of Occupational and Environmental Medicine, 46(4): 347-356, April 2004.
- 18. Siegel, M. "Involuntary Smoking in Restaurant Workplace: A Review of Employee Exposure and Health Effects." JAMA, 270:490-493, 1993.
- 19. Trout D, Decker J, Mueller C, Bernert JT, Pirkle J. Exposure of casino employees to environmental tobacco smoke. J Occup Environ Med. 1998 Mar;40(3):270-6.
- 21 Eisner Mark D. M.D. et al.: Bartender's Respiratory Health After Establishment of Smoke-free Bars and Tayerns. JAMA 1998 Dec 9:280(22):1909-1914
- 22. Mason-Dixon Polling & Research Inc. Survey of 625 registered Montana voters, May 23-25, 2005.